## Therapy of Degenerative Diseases of the Locomotorium

## J. Westphal, Sanatorium Landskrone

Bad Neuenahr

he referent gave an account of his experiences with the application of REGENERESEN over many years, acquired by more than ten thousand injections on about five hundred patients. Most of his patients are older than 65 years; 60 per cent of them are women.

The treatment always begins with an explanatory discussion with the patient to create an atmosphere of trust for the therapeutist and the natural therapies, and with a thorough examination, starting with non-invasive diagnostics, as an electrocardiogram, Doppler-sonography, and a test of the pulmonary function. These pre-examinations show three groups of diseases to be treated, either alone or in combination:

- metabolic diseases and circulatory disturbances
- peripheral and cerebral arterioscleroses
- diseases and ailments of the locomotorium (located mostly on knee and spine).


Sometimes these preliminary examinations show diseases, that were not known to the patient; for instance, cases of latent diabetes with glucose-levels of more than 200 mg per cent are frequently found.

The referent gave hints to other users of REGENERESEN how to explain the necessity and the advantage of a treatment with REGENERESEN to the patients. Such an explanation should not degress into biochemical details, but should show the effect of REGENERE-SEN on the activation of the organism's self-healing mechanisms, and on the improvement of metabolic and repairing processes in the cells, thus bettering or at least stopping progredient diseases.

The referent normally uses 20 ampoules of REGENERESEN together with other medicamental and physical therapies. With metabolic diseases, RN 13 is given together with REGENERESEN of stomach, large intestine, small intestine, lung, bronchi, vessel wall, and skin.

With diseases of the locomotorium, Osteochondrin S and REGENERESEN of osteoblasts, connective tissue, and musculature are always used, supplemented according to the ailment with vertebra, spinal cord, parathyroid gland, and bone marrow.

Many patients state at the follow-up examination, about three months after the preliminary examination, that they felt a real relief only two or three weeks before. This illustrates the fact, that especially with diseases of the locomo-



torium it can take two months or more, till a distinct success of the treatment is obtained. This success, however, then persists over a longer period (averagely for one to two years).